

- b) a naturally-occurring amino acid sequence having at least 90% sequence identity to the sequence of SEQ ID NO:1 or SEQ ID NO:3,
- c) a biologically-active fragment of the amino acid sequence of SEQ ID NO:1 or SEQ ID NO:3, and
- d) an immunogenic fragment of the amino acid sequence of SEQ ID NO:1 or SEQ ID NO:3.

44. **(Reiterated)** An isolated polypeptide of claim 43, having a sequence of SEQ ID NO:1 or SEQ ID NO:3.

---

45. **(Once amended)** An isolated polynucleotide encoding a polypeptide [of claim 43, or] selected from the group consisting of:

- C1
- a) an amino acid sequence of SEQ ID NO:1 or SEQ ID NO:3, and
  - b) a naturally-occurring amino acid sequence having at least 90% sequence identity to the sequence of SEQ ID NO:1 or SEQ ID NO:3,
- and a polynucleotide complementary thereto.
- 

46. **(Reiterated)** An isolated polynucleotide of claim 45, having a sequence of SEQ ID NO:2 or SEQ ID NO:4.

47. **(Reiterated)** A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 45.

48. **(Reiterated)** A cell transformed with a recombinant polynucleotide of claim 47.

C1  
C2

---

49. **(Once amended)** A method for producing a polypeptide [of claim 43] selected from the group consisting of:

- i) an amino acid sequence of SEQ ID NO:1 or SEQ ID NO:3,
- ii) a naturally-occurring amino acid sequence having at least 90% sequence identity to the sequence of SEQ ID NO:1 or SEQ ID NO:3,

iii) a biologically-active fragment of the amino acid sequence of SEQ ID NO:1 or SEQ ID NO:3, and

iv) an immunogenic fragment of the amino acid sequence of SEQ ID NO:1 or SEQ ID NO:3,

the method comprising:

a) culturing a cell of claim 48 under conditions suitable for expression of the polypeptide, [wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 43,] and

b) recovering the polypeptide so expressed.

50. **(Reiterated)** A transgenic organism comprising a polynucleotide of claim 47.

51. **(Reiterated)** An isolated antibody which specifically binds to a polypeptide of claim 43.

52. **(Reiterated)** An isolated polynucleotide comprising a sequence selected from the group consisting of:

- a) a polynucleotide sequence of SEQ ID NO:2 or SEQ ID NO:4,
- b) a naturally-occurring polynucleotide sequence having at least 90% sequence identity to the sequence of SEQ ID NO:2 or SEQ ID NO:4,
- c) a polynucleotide sequence complementary to a), and
- d) a polynucleotide sequence complementary to b).

53. **(Reiterated)** An isolated polynucleotide comprising at least 60 contiguous nucleic acids of a polynucleotide sequence of claim 52.

54. **(Reiterated)** A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 52, the method comprising:

- a) hybridizing the sample with a probe comprising at least 16 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide, and
- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

55. **(Reiterated)** A method of claim 54, wherein the probe comprises at least 30 contiguous nucleotides.

56. **(Reiterated)** A method of claim 54, wherein the probe comprises at least 60 contiguous nucleotides.

---

C3 57. **(Once amended)** A [pharmaceutical] composition comprising an effective amount of a polypeptide of claim 43 and a pharmaceutically acceptable excipient.

---

58. **(Reiterated)** A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 43, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 43 to a compound, and
- b) detecting agonist activity in the sample.

59. **(Reiterated)** A pharmaceutical composition comprising an agonist compound identified by a method of claim 58 and a pharmaceutically acceptable excipient.

60. **(Reiterated)** A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 43, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 43 to a compound, and
- b) detecting antagonist activity in the sample.

61. **(Reiterated)** A pharmaceutical composition comprising an antagonist compound identified by a method of claim 60 and a pharmaceutically acceptable excipient.

65. **(Reiterated)** A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 46, the method comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, and
- b) detecting altered expression of the target polynucleotide.

**Please add the following new claims:**

---

66. A method for assessing toxicity of a test compound, said method comprising:

- a) treating a biological sample containing nucleic acids with the test compound,
- b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 52 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 52,
- c) quantifying the amount of hybridization complex, and
- d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

67. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 52, the method comprising:

- a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and
- b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.